

## Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.



Information on the Microbiology of Eggs

by

Harry E. Goresline,  
Food Research Division,  
Bureau of Agricultural Chemistry and Engineering,  
U.S. Department of Agriculture,  
Washington, D.C.

The microbiology of eggs has received considerable study, since much of the spoilage and deterioration encountered in fresh, stored or processed eggs is due to the growth of microorganisms.

A certain percentage of fresh eggs contain bacteria at the time they are laid, but most freshly laid eggs are sterile. The white contains a natural bactericidal product known as lysozyme, which no doubt accounts for much of the freedom from contamination. The egg is, however, a medium in which bacterial multiplication will take place rapidly, and for this reason it is necessary that eggs be handled rapidly and under as sanitary conditions as possible if high quality material is to be expected. The presence of large numbers of microorganisms is an indication of improper handling, and often of unsanitary conditions. This applies to all phases of the egg industry, from the handling of the flock to the final use of the eggs.

Microbiology of Egg Handling and Storage

The care and storage of shell eggs is important from the microbiological standpoint, in that the type of eggs handled and the type of storage govern to a large extent the quality of the final product. If the flock is handled in such a way as produce a large percentage of "dirties" or the eggs are subjected to improper treatment a much higher loss from bacterial spoilage is to be expected.

It has been found that fresh dirty eggs with sound shells frequently become contaminated, probably through the large pores of the shell. The humidity of the air in storage is an important factor in the bacterial penetration of the shell of dirty and washed dirty eggs. If properly washed, dried and stored there will be comparatively little loss in storage.

Spoilage in storage is due chiefly to bacteria. Bacteria increase in numbers very rapidly at room temperature or higher, and for this reason it is essential to place the eggs in cool storage as soon as possible. The main types of bad eggs found during and after storage are green white, digested whites, and white and black rots. As a rule, these types show decided chemical decomposition and are heavily infected with bacteria.

The molding of eggs in storage gives them an unsightly appearance, but this trouble can be avoided by proper handling and control of humidity. Eggs which have been packed in wet cases or fillers are likely to show molding in a relatively short time. It has also

been observed that reinfection does occur when old infected cases are used.

### Microbiology of Egg Breaking

Rigid sanitary measures are essential in egg breaking plants if quality products are to be produced. Not only must the equipment be sterilized and the premises be sanitary in every respect, but care must be taken that bad eggs do not infect whole batches that have been broken out. For example, a large batch of good eggs can be made unfit for consumption by accidentally adding a musty egg through improper inspection. The musty odor and flavor in eggs infected with certain bacteria is very penetrating, and one musty egg may infect and scent an entire batch. The eggs should be broken separately, examined and smelled before adding to the main batch in the mixer. Bacteria increase in numbers by dividing into two individuals, and under ideal conditions this takes place every twenty to thirty minutes. It can be seen, therefore, why speed is essential in the handling of eggs and egg products and why contamination should be kept at a minimum. Freezing, drying or other processing should be done as rapidly as possible to keep down bacterial numbers, and thus prevent spoilage or loss of quality. Breaking equipment should be sterilized after a bad egg has come in contact with it, otherwise contamination will spread to eggs of good quality. The temperature of freezing storage should be kept low enough to keep the product solidly frozen at all times.

The main thing in the handling of eggs, from the bacteriological standpoint, is that to maintain quality, appearance and wholesomeness the hens should be properly cared for, and the eggs should be handled, stored or broken out under the best sanitary conditions. Proper temperature of storage, proper humidity and careful methods of handling will eliminate most of the trouble due to bacterial spoilage and deterioration.

A bulletin is now being prepared by the Department of Agriculture covering the various phases of egg production, storage, processing, and utilization.

There is attached a bibliography on the microbiology of eggs which may prove helpful.

The Microbiology of Eggs\*

A Selected Bibliography

Compiled by

Helen F. Smart,

Food Research Division,

Bureau of Agricultural Chemistry and Engineering,

U.S. Department of Agriculture,

Washington, D. C.

1. Microbiology of Shell Eggs:

Lamson, G. H.

1909. Infection and Preservation of Eggs. Bull. 55, pp. 203-14,  
Conn. Agr. Exp. Sta. (Storrs).

Pennington, M. E.

1910. A Chemical and Bacteriological Study of Fresh Eggs. Jour.  
Biol. Chem., vol. 7, pp. 109-32.

Maurer, O.

1911. Bacteriological Studies on Eggs. Bull. 180, Kans. State Agr.  
Exp. Sta. (Manhattan).

Stiles, G. W., and Bates, C.

1912. A Bacteriological Study of Shell, Frozen and Desiccated Eggs  
Made under Laboratory Conditions at Washington. Chem. Bull.  
158, U. S. Dept. Agr.

Rettger, L. F.

1913. The Bacteriology of the Hen's Egg with Special Reference to  
its Freedom from Microbic Invasion. Bull. 75, Conn. Agr. Exp.  
Sta. (Storrs).

Bushnell, L. D., and Maurer, O.

1914. Some Factors Influencing the Bacterial Content and Keeping  
Quality of Eggs. Bull. 201, Kans. State Agr. Exp. Sta. (Man-  
hattan).

Pennington, M. E., Jenkins, M. K., St. John, E. Q., and Hicks, W. B.

1914. A Bacteriological and Chemical Study of Commercial Eggs in  
the Producing Districts of the Central West. Dept. Bull. 51, U.  
S. Dept. Agr.

Hadley, P. B., and Caldwell, D. W.

1916. The Bacterial Infection of Eggs. Bull. 164, Rhode Island Agr.  
Exp. Sta. (Kingston).

Tanner, F. W.

1919. Bacteriology and Mycology of Foods. John Wiley & Sons, Inc.,  
New York, N. Y., 592 pp.

---

\*These books should be available for consultation through comprehensive  
public or technical libraries.



Tissier, H.

1926. Microbes de la putréfaction des Oeuf. Compt. Rend. Soc. Biol., vol. 94, no. 7, pp. 446-47. Abst. - Biol. Abst., vol. 1, no. 4, p. 510 (1927)

Turner, A. W.

1927. *Achromobacter perolens* (n. sp.) the Cause of Mustiness in Eggs. Australian Jour. Exp. Biol. & Sci., vol. 4, part 1, pp. 57-60.

Bohart, R. M.

1930. Anaerobic Bacteria Causing Black Rot of Eggs. American Jour. Hyg., vol. 11, pp. 168-73.

Spanswick, M. P.

1930. Cause of Mustiness in Eggs. Am. Jour. Pub. Health, vol. 20, no. 1, pp. 73-74.

Levine, M., and Anderson, D.

1930. Some Characteristics of Bacteria Causing Mustiness in Eggs. Jour. Bac., vol. 19, p. 55.

James, L. H., and Swenson, T. L.

1930. A study of Pin Spot Molding on Shell Eggs in Cold Storage. U. S. Egg and Poultry Mag., vol. 25, no. 12, pp. 47, 60.

Andresen, F.

1932. The Occurrence of Bacteria in Eggs. Inaug. Diss. Tierarztl. Hochsch. (Berlin), vol. III, p. 41.

Levine, M., and Anderson, D.

1932. Two New Species of Bacteria Causing Mustiness in Eggs. Jour. Bact., vol. 23, pp. 337-47.

Bryant, R. L., and Sharp, P. F.

1934. Effect of Washing on the Keeping Quality of Hen's Eggs. Jour. Agr. Res., vol. 48, p. 67.

Pennington, M. E.

1937. Cladosporia, A Study of Dust Storms and Mold on Eggs. U. S. Egg and Poultry Mag., vol. 43, no. 1, p. 24.

Miles, A. A., and Halnan, E. T.

1937. A New Species of Microorganism Causing Black Rot in Eggs. Jour. Hyg. (London), vol. 37, no. 1, p. 79.

Sharp, P. F.

1937. Preservation and Storage of Hen's Eggs. Food Research, vol. 2, no. 6, pp. 477-98.

Swenson, T. L.

1938. The Storage of Shell Eggs. Food Research, vol. 3, no. 6, pp. 599-608.

Haines, R. B.

1933. Observations on the Bacterial Flora of the Hen's Egg, with A Description of New Species of *Proteus* and *Pseudomonas* Causing Rots in Eggs. Jour. Hyg. (British), vol. 38, p. 338.

Haines, R. B.

1939. Microbiology in the Preservation of the Hen's Egg. British Food Inv. Bd., Spec. Rpt. 47, His Majesty's Stationery Office, London.

Turner, A. W., and Arnold L.

1939. Factors Involved in the Microbic Invasion of Hen's Eggs. U. S. Egg and Poultry Mag., vol. 45, no. 10, p. 599.

## 2. Effects of Cleaning and Storage Methods on the Keeping Quality of Eggs:

Jenkins, M. K., and Swan, C.

1920. Effects of Cold Storage on Shell Eggs. Ice and Refrig., vol. 58, no. 3, pp. 140-47.

Sharp, P. F., and Stewart, G. F.

1931. Carbon Dioxide and the Keeping Quality of Eggs. U. S. Egg and Poultry Mag., vol. 37, no. 6, pp. 30-32, 63-65.

Bryant, R. L., and Sharp, P. F.

1934. Effect of Washing on the Keeping Quality of Hens' Eggs. Jour. Agr. Res., vol. 48, p. 67.

McCammon, R. B., Pittman, M. S., and Wilhelm, L. A.

1934. The Odor and Flavor of Eggs. Poultry Sci., vol. 13, no. 2.

Sharp, P. F., and Stewart, G. F.

1936. Effect of Relative Humidity on the Growth of Mold on Eggs in Storage. Cornell Univ. Agr. Exp. Sta. Memoir 191.

Sharp, P. F., Stewart, G. F., and Huttar, J. C.

1936. Effect of Packing Material on the Flavor of Storage Eggs. Cornell Univ. Memoir 198.

Moran, T.

1937. Gas Storage of Eggs. Jour. Soc. Chem. and Ind. (London), vol. 56, p. 967.

Michael, K. E., and Mallmann, W. L.

1938. Mold Formation on Egg Packages. U. S. Egg and Poultry Mag., vol. 44, no. 5, pp. 293, 317.

Swenson, T. L.

1938. Storage of Shell Eggs. Food Research, vol. 3, no. 6, pp. 599-608.

Funk, E. M.

1938. Improving the Keeping Quality of Eggs by Cleaning with Sodium Hydroxide. Res. Bull. 277, Missouri Agr. Exp. Sta. (Columbia).

Ewell, A. W.

1938. Present Use and Future Prospects of Ozone in Food Control. Food Research, vol. 3, nos. 1, 2, pp. 101-108.

Moran, T.

1938. Gas Storage of Meat and Eggs. Food Research, vol. 3, nos. 1, 2, pp. 149-54.

Toop, E.

1939. The Occurrence of Mold on Egg Containers in Storage at A Chicago Plant. U. S. Egg and Poultry Mag., vol. 45, no. 7, pp. 397-99, 448.

Parsons, C. H., and Mink, L. D.

1939. A Review of Some Methods of Cleaning Eggs. U. S. Egg and Poultry Mag., vol. 45, no. 7, pp. 400, 433.

### 3. Microbiology of Frozen and Dried Eggs:

Stiles, G. W., and Bates, C.

1912. A Bacteriological Study of Shell, Frozen, and Desiccated Eggs. Chem. Bull. 158, U. S. Dept. Agr.

Pennington, M. E.

1912. Practical Suggestions for the Preparation of Frozen and Dried Eggs. Chem. Circ. 98, U. S. Dept. Agr.

Pennington, M. E., Jenkins, M. K., Stocking, W. A., Ross, S. H., St. John, E. Q., Hendrickson, M., and Hicks, W. B.

1916. A Study of the Preparation of Frozen and Dried Eggs in the Producing Section. Dept. Bull. 224, U. S. Dept. Agr.

Jenkins, M. K., and Pennington, M. E.

1918. The Installation and Equipment of an Egg-Breaking Plant. Dept. Bull. 663, U. S. Dept. Agr.

Tanner, F. W.

1919. Bacteriology and Mycology of Foods. John Wiley and Sons, Inc., New York, N. Y., 592 pp.

Redfield, H. W.

1920. Examination of Frozen Egg Products and Interpretation of Results. Dept. Bull. 846, U. S. Dept. Agr.

Jones, H. I., and Dubois, R.

1920. The Preservation of Eggs, Including A Bibliography on the Subject. Jour. Ind. Eng. Chem., vol. 12, no. 8, p. 751.

Thom, C., and Hunter, A. C.

1924. Hygienic Fundamentals of Food Handling. The Williams and Wilkins Co., Baltimore, Md., 228 pp.



Palmer, J. C.

1925. Report on the Methods for the Analysis of Dried Eggs. Jour. Assoc. Official Agr. Chem., vol. 8, p. 615.

De Bord, G. F.

1925. Effect of Dehydration upon the Bacterial Flora of Eggs. Jour. Agr. Res., vol. 31, no. 2, pp. 155-64.

Moran, T., and Piqué, J.

1926. The Storage of Eggs. Food Inv. Bd., Spec. Rept. 26, His Majesty's Stationery Office, London.

Bailey, M. E.

1932. A Study of the Gelation of Frozen Egg Magma. Thesis, Columbia Univ. Faculty of Pure Sci., New York, N. Y.

Swenson, T. L., and James, L. H.

1935. A Comparison Between Eggs Frozen at Zero Fahrenheit and Eggs Frozen at -109° Fahrenheit. U. S. Egg and Poultry Mag., vol. 41, no. 3, pp. 16-19.

Dack, G. M.

1937. Report, Public Health Section of Proceedings Poultry Products Conference (1935). Abst. - U. S. Egg and Poultry Mag., vol. 43, no. 11, pp. 698-99.

Sharp, P. F.

1937. Preservation and Storage of Hen's Eggs. Food Research, vol. 2, no. 6, pp. 477-98.

Sealey, J. O.

1939. Bacteria and Other Biological Factors in Relation to Frozen Foods. U. S. Egg and Poultry Mag., vol. 45, no. 6, pp. 355-57, 380.

Haines, R. B.

1939. Microbiology in the Preservation of the Hen's Egg. British Food Inv. Bd., Spec. Rept. 47, His Majesty's Stationery Office, London.

#### 4. The Lytic or Bactericidal Principle of Egg White:

Rettger, L. F., and Sperry, S. A.

1912. Antiseptic and Bactericidal Properties of Egg White. Jour. Med. Res., vol. 26, pp. 55-64.

Sperry, J. A.

1913. An Antiseptic and Bactericidal Property of Egg White. Science, vol. 38, p. 413.

Fleming, A., and Allison, V. D.

1924. On the Antibacterial Power of Egg White. *Lancet*, vol. I, pp. 1305-07.

Sandow, A.

1926. The Anti-Bacterial Activity of Egg White. *Proc. Soc. Exp. Biol. Med.*, Vol. 24, pp. 172-73.

Stark, C. N., and Sharp, P. F.

1927. The Relationship Between Hydrogen Ion Concentration of Egg White and the Growth of Anaerobes. *Abst. - Jour. Bact.*, vol. 13, no. 1, p. 59.

Sharp, P. F., and Whitaker, R.

1927. The Relation of pH Concentration of Egg White to its Germicidal Action. *Jour. Bact.*, vol. 14, pp. 17-46.

Sharp, P. F.

1929. The pH of the White as an Important Factor Influencing the Keeping Quality of Hen's Eggs. *Science*, vol. 69, no. 1784, pp. 278-80.

Meyer, K., Thompson, R., Palmer, J. W., and Khorazo, D.

1934. The Nature of Lysozyme Action. *Science*, vol. 79, no. 2038, p. 61.



